arXiv:math/0107113v1 [math.FA] 16 Jul 2001

ON LINEAR OPERATORS WITH p-NUCLEAR ADJOINTS

O.I.Reinov[†]

Abstract. If $p \in [1, +\infty]$ and T is a linear operator with p-nuclear adjoint from a Banach space X to a Banach space Y then if one of the spaces X^* or Y^{***} has the approximation property, then T belongs to the ideal N^p of operators which can be factored through diagonal oparators $l_{p'} \to l_1$. On the other hand, there is a Banach space W such that W^{**} has a basis and such that for each $p \in [1, +\infty], p \neq 2$, there exists an operator $T : W^{**} \to W$ with p-nuclear adjoint that is not in the ideal N^p, as an operator from W^{**} to W.

In a series of previous notes the author considered the question on the existence or nonexistence of non-p-nuclear operators with p-nuclear second adjoints in concrete Banach spaces. For example, in [3] such "bad" operators were constructed for every $p \in [1, 2)$ in spaces with Schauder bases.

Here we treat the analogous case for the operators *whose first adjoints are p-nuclear*. Already in the paper [2, §4] the corresponding assertions were formulated (without proofs) in which the spaces under consideration, however, did not possesses bases, — in the best case only one of the spaces, in which the operators acted, had the approximation property, but yet not bounded AP. In this note, proving partly those assertions from [2], we show that the similar examples can be found in spaces with bases (and even in better spaces). Simultaneously, some sufficient conditions are given for the positive answer to the corresponding question. These questons are connected with the p-nuclearity of the adjoint operator (moreover, those conditions are shown to be close to the necessary ones).

All the spaces under consideration are Banach spaces. We usually denote the elements of the spaces by the corresponding small letters: $x \in X$, $y \in Y, \dots,$ $x' \in$

[†] This work was done with partial support of the Ministry of the general and professional education of Russia (Grant 97-0-1.7-36) and FCP "Integracija", reg. No. 326.53.

[‡] AMS Subject Classification: 47B10. Hilbert–Schmidt operators, trace class operators, nuclear operators, p-summing operators, etc.

Key words: p-nuclear operators, dual ideal, bases, approximation properties, tensor products.

Typeset by $\mathcal{AMS}$-TEX

X^*, $y'' \in Y^{**}, \dots$. For $p \in [1, +\infty]$, the conjugate exponent p' is defined by the relation $1/p + 1/p' = 1$. By $L(X, Y)$ it is denoted the space of all (linear continuous) operators from X to Y with its standard norm. We shall consider every Banach space also as a subspace in its second dual, usually without introducing any additional notations for the natural imbedding $X \to X^{**}$. However, in the case of a necessity, we will denote by π_X this canonical imbedding.

Let us recall that *a Banach space has the* (*Grothendieck*) *approximation property* (the property AP), if the identity map on the space can be approximated, in the topology of compact convergence, by finite-dimensional operators.

For $p \in [1, \infty]$, an operator T, acting from X to Y, is said to be an *N^p-operator,* if it can be represented in the following form:

$$(1) \qquad Tx = \sum_{k=1}^{\infty} < x, x'_k, > \, y_k \qquad \text{for } x \in X,$$

where the sequences $\{x'_n\}_{n=1}^{\infty} \subset X^*$ and $\{y_n\}_{n=1}^{\infty} \subset Y$ are such that the quantity

$$(2) \qquad \alpha := \left(\sum_{k=1}^{\infty} \|y_k\|^p \right)^{1/p} \sup \left\{ \left(\sum_{k=1}^{\infty} |< x'_k, x >|^{p'} \right)^{1/p'} : \ x \in X, \ \|x\| \leq 1 \right\}$$

is finite (recall that p' is the conjugate exponent for p; in the case when one of the exponents p, p' is equal to infinity, one must understand the right part of the relation (2) properly). The set of all N^p-operators from X to Y is denoted by $N^p(X, Y)$, and the lowest bound $\inf \alpha$, where the *inf* is taken over all possible representations in the form (1) of the operator T, — by $\nu^p(T)$. For every $p \geq 1$ the class N^p of all N^p-operators is a Banach operator ideal [1]; for fixed spaces X, Y $N^p(X, Y)$ is a Banach space with the norm ν^p. Let us note that for the exponents $p, q \in [1, +\infty], p < q$, we always have: $N^p \subset N^q$ with the corresponding inequality for norms.

We denote by $X^* \widehat{\otimes}^p Y$ the tensor product associated with the space $N^p(X, Y)$: this is the completion of the algebraic tensor product $X^* \otimes Y$ via the norm ν_0^p, which is defined for $z \in X^* \otimes Y$ as the lowest bound, over all possible representations $z = \sum_{k=1}^{N} x'_k \otimes y_k$ in $X^* \otimes Y$, of the numbers

$$\left(\sum_{k=1}^{N} \|y_k\|^p \right)^{1/p} \sup \left\{ \left(\sum_{k=1}^{N} |< x'_k, x >|^{p'} \right)^{1/p'} : \ x \in X, \ \|x\| \leq 1 \right\}$$

It is not difficult to see that the space $N^p\,(X,Y)$ is the image of $X^*\widehat{\otimes}^p Y$ under the natural mapping $X^*\widehat{\otimes}^p Y \to L(X,Y)$. For $z \in X^*\widehat{\otimes}^p Y$, the induced operator we will denote by $\widetilde{z}$.

We shall need below the following reformulation of the corollary 1.2 from [2]:

$(*)$ There exists a reflexive separable Banach space E such that for each $r \neq 2$ the canonical mapping $E^*\widehat{\otimes}^r E \to N^r(E,E)$ is not one–to–one.

The dual space of $X^*\widehat{\otimes}^p Y$ is isometrically equal to the space $\Pi^d_{p'}(Y,X^{**})$, where by $\Pi^d_{p'}$ we denote the ideal, dual to the ideal of absolutely p'-summing operators with the corresponding norm (see [1]): if $z \in X^*\widehat{\otimes}^p Y$ and $U \in \Pi^d_{p'}(Y,X^{**})$, then the duality defined with the help of trace: trace $U \circ z$. Note that in case when one of the space X^* or Y possesses the AP, the canonical mapping $X^*\widehat{\otimes}^p Y \to L(X,Y)$ is one–to–one and thus we can write in this case: $X^*\widehat{\otimes}^p Y = N^p\,(X,Y)$.

The assertion $(*)$ can be reformulated by the following way:

$(**)$ There exists a reflexive separable Banach space E such that for each $r \neq 2$ one can find a tensor $z \in E^*\widehat{\otimes}^r E$ and an operator $U \in \Pi^d_{r'}(E,E)$ for which trace $U \circ z = 1$ and the associated with z operator $\widetilde{z} = 0$.

Theorem 1. *Let $p \in [1,+\infty]$, $T \in L(X,Y)$ and either $X^* \in AP$ or $Y^{***} \in AP$. If $T \in N^p(X,Y^{**})$, then $T \in N^p(X,Y)$. In other words, under these conditions from the p-nuclearity of the conjugate operator T^* it follows that the operator T belongs to the space $N^p(X,Y)$.*

Proof. Suppose there exists such an operator $T \in L(X,Y)$, that $T \notin N^p(X,Y)$, but $\pi_Y\,T \in N^p(X,Y^{**})$. Since either X^* or Y^{**} has the AP, $N^p(X,Y^{**}) = X^*\widehat{\otimes}^p Y^{**}$. Therefore the operator $\pi_Y\,T$ can be identified with the tensor element $t \in X^*\widehat{\otimes}^p Y^{**}$; in addition, by the choice of T, $t \notin X^*\widehat{\otimes}^p Y$ (the space $X^*\widehat{\otimes}^p Y$ is considered as a subspace of the space $X^*\widehat{\otimes}^p Y^{**}$). Hence there is an operator $U \in \Pi^d_{p'}(Y^{**},X^{**}) = \left(X^*\widehat{\otimes}^p Y^{**}\right)^*$, with the property that trace $U \circ t =$ trace $(t^* \circ (U^*|_{X^*})) = 1$ and trace $U \circ \pi_Y \circ z = 0$ for each $z \in X^*\widehat{\otimes}^p Y$. ¿From the last it is follows that, in particular, $U\pi_Y = 0$ and $\pi_Y^*\,U^*|_{X^*} = 0$. In fact, if $x' \in X^*$ and $y \in Y$, then

$$< U\pi_Y\,y, x' >=< y, \pi_Y^*\,U^*|_{X^*}x' >= \text{trace }\, U \circ (x' \otimes \pi_Y(y)) = 0.$$

Evidently, the tensor element $U \circ t$ induces the operator $U\pi_Y T$, which is equal identically to zero.

If $X^* \in AP$ then $X^* \widehat{\otimes}^p Y^{**} = N^p(X, Y^{**})$ and, therefore, this tensor element is zero what is contradicted to the equality trace $U \circ t = 1$.

Let now $Y^{***} \in AP$. In this case

$$V := (U^*|_{X^*}) \circ T^* \circ \pi_Y^* : \; Y^{***} \to Y^* \to X^* \to Y^{***}$$

uniquely determines a tensor element t_0 from the projective tensor product $Y^{****} \widehat{\otimes} Y^{***}$. Let us take any representation $t = \sum x_n' \otimes y_n''$ for t as an element of the space $X^* \widehat{\otimes}^p Y^{**}$. Denoting for the brevity the operator $U^*|_{X^*}$ by U_*, we obtain:

$$\begin{aligned} V y''' &= U_* \, (T^* \pi_Y^* \, y''') = U_* \, ((T^* \pi_Y^* \pi_{Y^*}) \, \pi_Y^* \, y''') = U_* \, ((\pi_Y T)^* \, \pi_{Y^*}) \, \pi_Y^* \, y''') = \\ &= U_* \left((\sum y_n'' \otimes x_n') \, \pi_{Y^*}) \, \pi_Y^* \, y''' \right) = U_* \left(\sum < y_n'', \pi_Y^* \, y''' > \, x_n' \right) = \\ &= \sum < \pi_Y^{**} y_n'', y''' > \, U_* x_n'. \end{aligned}$$

So, the operator V (or the element t_0) has in the space $Y^{****} \widehat{\otimes} Y^{***}$ the representation

$$V = \sum \pi_Y^{**}(y_n'') \otimes U_*(x_n').$$

Therefore,

$$\text{trace } t_0 = \text{trace } V = \sum < \pi_Y^{**}(y_n''), U_*(x_n') >= \sum < y_n'', \pi_Y^* \, U_* x_n' >= \sum 0 = 0.$$

On the other hand,

$$V y''' = U_* \, (\pi_Y T)^* \, y''' = U_* \circ t^*(y''') = U_* \left(\sum < y_n'', y''' > \, x_n' \right) = \sum < y_n'', y''' > \, U_* x_n',$$

whence $V = \sum y_n'' \otimes U_*(x_n')$. Therefore

$$\text{trace } t_0 = \text{trace } V = \sum < y_n'', U_* x_n' >= \sum < U y_n'', x_n' >= \text{trace } U \circ t = 1.$$

The obtained contradiction completes the proof of the theorem. ■

Theorem 2. *For each $r \in [1, \infty], r \neq 2$, there exist a separable space W and an operator $T \in L(W^{**}, W)$ such that W^{**} has a basis, $T \in N^r(W^{**}, W^{**})$, but $T \notin N^r(W^{**}, W)$.*

Proof. Let us fix $r \in [1, \infty], r \neq 2$ and take the triple (E, z, U) from the assertion $(**)$. Let W be a separable space such that W^{**} has a basis and there exists a linear homomorphism φ from W^{**} to E with the kernel $W \subset W^{**}$ so that the subspace $\varphi^*(E^*)$ in complemented in W^{***} (see [4]). Lift the tensor element z, lying in $E^*\widehat{\otimes}^r E$, up to an element[1] $\alpha \in E^*\widehat{\otimes}^r W^{**}$, so that $\varphi \circ \alpha = z$, and set $V := U \circ \varphi$. Since trace $V \circ \alpha =$ trace $U \circ z = 1$ and W^{**} has the AP, then $\widetilde{\alpha} = \alpha \neq 0$. Besides, the operator $\widetilde{\varphi \circ \alpha} : E \to W^{**} \to E$, associated with the tensor $\varphi \circ \alpha$, is equal to zero. Therefore $\alpha(E) \subset \operatorname{Ker} \varphi = W \subset W^{**}$, that is the operator α is acted from E into W.

Since the subspace $\varphi^*(E^*)$ is complemented in W^{***}, then $\alpha \circ \varphi \in W^{***}\widehat{\otimes}^r W = N^r(W^{**}, W)$ iff $\alpha \in E^*\widehat{\otimes}^r W = N^r(E, W)$.

If $\alpha \in N^r(E, W)$, then for its arbitrary (nonzero!) N^p-representation of the form $\alpha = \sum e'_n \otimes w_n$ the composition $\varphi \circ \alpha$ is a zero tensor element in $E^*\widehat{\otimes} E$; but this composition represents the element z, which, by its choice, can not be zero one. Thus, $\alpha \notin N^r(E, W)$ and, thereby, $\alpha \circ \varphi \notin W^{***}\widehat{\otimes}^r W = N^r(W^{**}, W)$. On the other hand, certainly, $\alpha \circ \varphi \in W^{***}\widehat{\otimes}^r W^{**} = N^r(W^{**}, W^{**})$. ■

So, the approximation conditions imposed on X and Y in the theorem 1 are essential.

It is clear that from $(**)$ and from the proof of Theorem 2 it follows that there exists a Banach space W, for which the conclusion of the theorem is valid for all mentioned values of the parameter r.

REFERENCES

1. A. Pietsch, *Operator ideals*, North-Holland (1978), 536 p.
2. O. I. Reinov, *Approximation properties of order p and the existence of non-p-nuclear operators with p-nuclear second adjoints*, Math. Nachr. **109** (1982), 125-134.
3. O. I. Reinov, *Approximation properties* AP_s *and p-nuclear operators* (*the case when* $0 < s \leq 1$), Zapiski nauchn. sem. POMI **270** (2000), 277-291 (in Russian).
4. J. Lindenstrauss, *On James' paper "Separable Conjugate Spaces"*, Israel J. Math. **9** (1971), 279-284.

Oleg I. Reinov
Department of Mathematics

---

[1] If $z = \sum_{k=1}^{\infty} e'_k \otimes e_k$ is any representation of z in $E^*\widehat{\otimes}^r E$, then we take $\{w''_n\} \subset W^{**}$ in such a way that the last sequence is absolutely r-summing and $\varphi(w''_n) = e_n$ for every n.

St Petersburg University
St Peterhof, Bibliotech pl 2
198904 St Petersburg, Russia
E-mail address: `orein@orein.usr.pu.ru`